**Docket #:** S19-557

# Continuous, Real-Time Monitoring of Biomolecules in Live Animals

Researchers at Stanford have developed methods for the continuous real-time monitoring of biomolecules in live subjects.

Real-time biosensors that can continuously measure circulating biomolecules in vivo would revolutionize the healthcare system by enabling truly personalized medicine – providing valuable insights into a patient's health status and their response to therapeutics, allowing clinicians to tailor therapeutic regimens to consistently deliver maximum efficacy with minimal side effects. Unfortunately, current real-time biosensors are limited to a handful of analytes and low sensitivity. Further, with current technologies, understanding individualized therapeutic response kinetics is laborious and most treatment decisions are made with a combination of representative population data combined with trial and error.

#### Stage of Research

The inventors have developed methods for the continuous and simultaneous monitoring of multiple analytes with picomolar sensitivity and sub-second temporal resolution. Their methods and microfluidic device function together as a biosensor which constantly processes samples and integrates several steps, such as purification, incubation and washing. The inventors demonstrate real-time continuous detection of glucose and insulin in live diabetic rats using a fluorescence sandwich immunoassay. Their device can measure transient changes in glucose and insulin levels. Further, they demonstrate the capacity to resolve inter-individual differences in pharmacokinetic responses to therapeutics.

## **Applications**

• Real-time in vivo detection and concentration measurements of multiple analytes using sensors within microfluidic devices

 Treating a disease or disorder via continuous or intermittent administration of a therapeutic based on determined concentration(s) of one or more analytes associated with the disease or disorder

### **Advantages**

- Sensor exhibits sufficient sensitivity, specificity, and dynamic range in measuring low abundance molecules directly in whole blood to resolve temporal fluctuations over short timescales
- Device incubation time of 30 seconds, drastic improvement to traditional ELISAbased detection technologies
- Core components can be applied for in vivo, multiplexed, and continuous measurement of any target molecule of interest

#### **Publications**

Poudineh M, Maikawa CL, Ma EY, Pan J, Mamerow D, Hang Y, Baker SW,
Berirami A, Yoshikawa A, Eisenstein M, Kim S, Vuckovic J, Appell EA, Soh HT. <u>A</u>
<u>fluorescence sandwich immunoassay for the real-time continuous detection of glucose and insulin in live animals. Nat Biomed Eng.</u> 2021.

#### **Patents**

Published Application: WO2021146612
Published Application: 20230061218

#### **Innovators**

- H. Tom Soh
- Mahla Poudineh
- Jing Pan

## **Licensing Contact**

## **Kimberly Griffin**

Technology Licensing and Strategic Alliances Manager

<u>Email</u>